

Separable Faces and Renormings of Non-separable Banach Spaces.

Nine Open Problems since 1975-76, 2007 and 2020.

José Orihuela

Universidad de Murcia

We analyze question 18 of J. Lindenstrauss in [5]. We prove that a Banach space E with a norming subspace $F \subset E^*$ has an equivalent $\sigma(E, F)$ -lower semicontinuous LUR norm if, and only if, there is a sequence $\{A_n : n = 1, 2, \dots\}$ of subsets of E such that, given any $x \in E$ and $\epsilon > 0$, there is a $\sigma(E, F)$ -open half-space H and $p \in \mathbb{N}$ such that $x \in H \cap A_p$ and the slice $H \cap A_p$ can be covered with countable many sets of diameter less than ϵ . Thus E has an equivalent $\sigma(E, F)$ -lower semicontinuous LUR norm if, and only if, it has another one with separable denting faces,[9, 10] **This result completely solves four problems asked in [7, Question 6.33, p.128] extending Troyanski's fundamental results (see Chapter IV in [1]), and others ones in [3, 6]. Moreover, LUR renormings are possible at points of separable faces wich could be glued as a σ -slicely isolated family of faces [7], of the unit sphere of E .** Among new examples covered by this results are Banach spaces $C(K)$, where K is a Rosenthal compact space $K \subset \mathbb{R}^\Gamma$ i.e., a compact space of Baire one functions on a Polish space Γ , with at most countably many discontinuity points for every $s \in K$, **which solves three problems asked in [7, Question 6.23, p.125].** Previously, it was only known for K being separable too, see [4] where the σ -fragmentability of $C(K)$ was already proved for non separable K , and a conjecture for the pointwise lower semicontinuous and LUR renorming presented here was posed, details will appear in [8].

For strictly convex renormings **we solve a recent question of R. Smith [11] giving a final answer to Lindenstauss question 18 in [5], see [2].**

Joint work with V. Montesinos. Research partially supported by Project PID2021-122126NB-C33 and Fundación Séneca - ACyT Región de Murcia project 21955/PI/22.

References

- [1] R. Deville, G. Godefroy, and V. Zizler. *Smoothness and renormings in Banach spaces*, volume 64 of *Pitman Monographs and Surveys in Pure and Applied Mathematics*. Longman Scientific & Technical, Harlow, 1993.
- [2] S. Ferrari, V. Montesinos and J. Orihuela. A separation method through convex sets for strictly convex renormings of Banach spaces. Preprint 2023.
- [3] F. García, L. Oncina, J. Orihuela, and S. Troyanski. Kuratowski's index of non-compactness and renorming in Banach spaces. *J. Convex Anal.*, 11(2):477–494, 2004.
- [4] R. Haydon, A. Moltó, and J. Orihuela. Spaces of functions with countably many discontinuities. *Israel J. Math.*, 158:19–39, 2007.
- [5] J. Lindenstrauss. Some open problems in Banach space theory. In *Séminaire Choquet. Initiation à l'analyse tome 15, 1975-1976, Exp. No. 18*, pages 1–9. Secrétariat mathématique, Paris, 1975-76.
- [6] A. Moltó, J. Orihuela, and S. Troyanski. Locally uniformly rotund renorming and fragmentability. *Proc. London Math. Soc. (3)*, 75(3):619–640, 1997.
- [7] A. Moltó, J. Orihuela, S. Troyanski, and M. Valdivia. *A nonlinear transfer technique for renorming*, volume 1951 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2009.
- [8] A. Moltó, V. Montesinos, J. Orihuela and S. Troyanski. Rosenthal compact spaces and renormings. Work in progress.
- [9] V. Montesinos and J. Orihuela. Separable slicing and locally uniformly rotund renormings of Banach spaces. Submitted January 2023
- [10] V. Montesinos and J. Orihuela. Weak compactness and separability of faces for convex renormings of Banach spaces. Preprint 2023
- [11] R.J. Smith A topological characterization of dual strict convexity in Asplund spaces. *J. Math. Anal. Appl.* 485 (2020) 123806